



# Difference Equations and Chaos

10-16-2006





# Opening Discussion


- What did we talk about last class?
- Do you have any questions about the project?





# Difference Equations

- Similar to differential equations are difference equations. These are discrete equations where we calculate the next value of the system from the previous one.
- These systems are sometimes referred to as mappings.
- Our numerical solutions to differential equations actually convert them to difference equations.





# The Logistic Map

- A common example of a map is the logistic map. The formula for the logistic map is extremely simple:  $x_{n+1} = rx_n(1-x_n)$ .
- Iterating a simple 1-D map like this can be viewed by drawing a “cobweb diagram”. The formula is quadratic in  $x$ , but opens downward.
- Fixed points are places where the curve crosses the  $y=x$  line. Depending on the slope at that the intersection the fixed points might be stable or unstable.
- Let's see if we can write code to draw a cobweb diagram.



- If we vary the value of  $r$ , interesting things happen to the behavior of this system. At small values there is a single, stable fixed point. At larger values, that fixed point becomes unstable and we get a period-2 cycle instead. This split is called a bifurcation.
- Increasing the value a bit more produces another bifurcation to a period-4 cycle.
- The separation between bifurcations gets smaller and the system actually becomes chaotic.
- Let's write code to draw a bifurcation diagram.



- The midterm will be tomorrow. It is open book and open Matlab.

